

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human MSP/MST1 in direct ELISAs and Western blots. In direct ELISAs and Western blots, less than 5% cross-reactivity with recombinant human HGF is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant human MSP/MST1 Gln19-Gly711 Accession # AAA59872
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Neutralization	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Macrophage stimulating protein (MSP), also known as HGF-like protein, and scatter factor-2, is a member of the HGF family of growth factors (1). MSP is secreted as an inactive single chain precursor (pro-MSP) that contains a PAN/APPLE-like domain, four kringle domains, and a peptidase S1 domain which lacks enzymatic activity (2). Human MSP shares 79% aa sequence identity with mouse MSP and 44% aa sequence identity with human HGF. Pro-MSP is secreted by hepatocytes under the positive and negative control of CBP in complex with either HNF-4 or RAR, respectively (3). Circulating pro-MSP is proteolytically cleaved in response to tissue injury to yield biologically active disulfide linked heterodimers consisting of a 45-62 kDa alpha and a 25-35 kDa beta chain (4, 5). Pro-MSP can be activated by MT-SP1, a transmembrane protease that is expressed on macrophages and is upregulated in many cancers (6). Heterodimeric MSP as well as the isolated beta chain bind to MSP R/Ron with high-affinity, although only heterodimeric MSP can induce receptor dimerization and signaling (7, 8). MSP induces macrophage and keratinocyte proliferation and osteoclast activation (9, 10). It also inhibits LPS- or IFN-induced iNOS and IL-12 expression by macrophages and prevents apoptosis of epithelial cells separated from the ECM (11, 12).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.