

DESCRIPTION

Species Reactivity	Human/Mouse
Specificity	Detects human and mouse BMP-7 in Western blots. In this format, less than 1% cross-reactivity with recombinant human (rh) BMP-2, rhBMP-3, rhBMP-4, rhBMP-5, and rhBMP-6 is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant human BMP-7 Ser293-His431 Accession # P18075
Conjugate	Alexa Fluor 405 Excitation Wavelength: 405 nm Emission Wavelength: 421 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Bone morphogenetic protein 7 (BMP-7), also known as osteogenic protein 1 (OP-1), is a widely expressed TGF-β superfamily member with important functions during embryogenesis, in the adult, and in disease (1, 2). Human BMP-7 is synthesized with a 29 amino acid (aa) signal sequence, a 263 aa propeptide, and a 139 aa growth factor domain (3, 4). The growth factor domain of human BMP-7 shares 98% aa sequence identity with mouse and rat BMP-7. The BMP-7 propeptide is cleaved intracellularly but often remains associated with the mature C-terminus. Based on in vivo and in vitro studies, BMP-7 has the potential to be secreted as a disulfide-linked mature homodimer, or particularly as a heteromeric complex that consists of two propeptides noncovalently associated with a mature disulfide-linked homodimer (5, 6). The presence of the propeptides in BMP-7 appears to stabilize the molecule and provide a docking mechanism for extracellular storage on molecules such as fibrillin-1 and -2 (5, 6). The propeptides themselves do not impart latency to the complex. BMP-7 binding to type II receptors rapidly displaces the prodomain:mature molecule interaction and has no effect on activity. But it is suggested that immobilized BMP-7 (via prodomain:fibrillin) is inactive, allowing for possible long-term storage of the molecule (6). BMP-7 interacts with the type 2 receptors Activin RIIA, Activin RIIIB, and BMPRII and the type 1 receptors Activin RIA, BMPRI-IA, and BMPRI-IB (2, 6). BMP-7 may also be processed into a disulfide-linked heterodimer with either BMP-2 or BMP-4. Such complexes may show increased potency and range of activity compared to BMP-7 homodimers (7 - 9). BMP-7 plays a role in a variety of organ systems. It promotes new bone formation and nephron development (10, 11), inhibits the branching of prostate epithelium (12), and antagonizes epithelial-mesenchymal transition (EMT) (13 - 15). In pathological conditions, BMP-7 inhibits tumor growth and metastasis (14), ameliorates fibrotic damage in nephritis (13), and promotes neuroregeneration following brain ischemia (16).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.