

Antigen Affinity-purified Polyclonal Goat IgG Catalog Number: AF3630

DESCRIPTION

| DECONTRA TION | |
|--------------------|--|
| Species Reactivity | Human |
| Specificity | Detects human RGM-B in direct ELISAs and Western blots. In direct ELISAs and Western blots, approximately 50% cross-reactivity with recombinant mouse RGM-B is observed and less than 1% cross-reactivity with recombinant human (rh) RGM-A and rhRGM-C is observed. |
| Source | Polyclonal Goat IgG |
| Purification | Antigen Affinity-purified |
| Immunogen | Mouse myeloma cell line NS0-derived recombinant human RGM-B isoform 1 Gly46-SerSer411 Accession # Q6NW40 |
| Formulation | Lyophilized from a 0.2 μm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 μm filtered solution in PBS. |

APPLICATIONS Relase Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website. Recommended Concentration Sample Western Blot 0.1 µg/mL Recombinant Human RGM-B (Catalog # 3630-RG) Flow Cytometry 0.25 µg/10⁶ cells See Below CyTOF-ready Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.

DATA



Detection of RGM-B in A172 Human Cell Line by Flow Cytometry. A172 human glioblastoma cell line was stained with Goat Anti-Human RGM-B Antigen Affinity-purified Polyclonal Antibody (Catalog # AF3630, filled histogram) or isotype control antibody (Catalog # Catalog # AB-108-C, open histogram), followed by Allophycocyanin-conjugated Anti-Goat IgG Secondary Antibody (Catalog # Catalog # F0108). View our protocol for Staining Membrane-associated Proteins.

| PREPARATION AND STORAGE | |
|-------------------------|--|
| Reconstitution | Reconstitute at 0.2 mg/mL in sterile PBS. |
| Shipping | The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C |
| Stability & Storage | Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 °C as supplied. 1 month, 2 to 8 °C under sterile conditions after reconstitution. 6 months, -20 to -70 °C under sterile conditions after reconstitution. |

Rev. 6/3/2022 Page 1 of 2



Global bio-techne.com info@bio-techne.com techsupport@bio-techne.com TEL +1 612 379 2956 USA TEL 800 343 7475 Canada TEL 855 668 8722 China TEL +86 (21) 52380373 Europe | Middle East | Africa TEL +44 (0)1235 529449



Human RGM-B Antibody

Antigen Affinity-purified Polyclonal Goat IgG Catalog Number: AF3630

BACKGROUND

RGM-B, also known as DRAGON, is a 40 kDa member of the repulsive guidance molecule (RGM) family of GPI-linked neuronal and muscle membrane proteins (1, 2). It is synthesized as a preproprotein that consists of a 45 amino acid (aa) signal sequence, a 368 aa mature region, and a 24 aa C-terminal prosegment (3). RGM-B contains an RGD motif, two potential N-linked glycosylation sites, and an abbreviated von Willebrand factor domain. There is a potential proteolytic cleavage site within the VWF domain (4). Alternative splicing may generate isoforms of RGM-B with N-terminal extensions or truncation following the VWF domain. Mature human RGM-B shares 52% and 36% aa sequence identity with the comparable regions of RGM-A and RGM-C, respectively. It shares 98%, 92%, 92%, and 78% aa sequence identity with macaque, mouse, bovine, and chicken RGM-B, respectively. RGM-B is expressed in the developing and adult nervous system, particularly in the dorsal root ganglia and mantle layer of the spinal cord (3-5). In mouse, it shows a complementary, non-overlapping distribution with RGM-A (2-5). RGM-B is also expressed in fetal and adult enteric ganglia and in postnatal intestinal epithelium (6). RGM-B expression has been detected in neuronal cell bodies and proximal axonal segments (4) but is also present on the cell surface, where it interacts homophilically and mediates neuronal adhesion (3). RGM-B additionally functions as a BMP coreceptor. It directly binds BMP-2 and -4 but not other TGF-β family proteins (7). RGM-B associates with BMP type I (ALK-2, -3, -6) and type II (Activin RIIA, Activin RIIB) receptors and enhances BMP signaling (7).

References:

- 1. Monnier, P.P. et al. (2002) Nature 419:392.
- 2. Schmidtmer, J. and D. Engelkamp (2004) Gene Exp. Patterns 4:105.
- 3. Samad, T.A. et al. (2004) J. Neurosci. 24:2027.
- 4. Niederkofler, V. et al. (2004) J. Neurosci. 24:808.
- 5. Oldekamp, J. et al. (2004) Gene Exp. Patterns 4:283.
- 6. Metzger, M. et al. (2005) Dev. Dyn. 234:169.
- 7. Samad, T.A. et al. (2005) J. Biol. Chem. 280:14122.

Rev. 6/3/2022 Page 2 of 2



Global bio-techne.com info@bio-techne.com techsupport@bio-techne.com TEL +1 612 379 2956 USA TEL 800 343 7475 Canada TEL 855 668 8722 China TEL +86 (21) 52380373 Europe | Middle East | Africa TEL +44 (0)1235 529449