

DESCRIPTION


Species Reactivity	Human/Mouse/Rat
Specificity	Detects human, mouse, and rat HSP90 in Western blots. Detects recombinant human HSP90 α and HS90 β in Western blots.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human HSP90 α Met1-Asp732 Accession # P07900
Formulation	Lyophilized from a 0.2 μ m filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 μ m filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	0.5 μ g/mL	See Below

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 $^{\circ}$ C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> • 12 months from date of receipt, -20 to -70 $^{\circ}$C as supplied. • 1 month, 2 to 8 $^{\circ}$C under sterile conditions after reconstitution. • 6 months, -20 to -70 $^{\circ}$C under sterile conditions after reconstitution.

BACKGROUND

The heat shock protein-90 kDa (HSP90) is a composite name for a large group of genes whose molecular weights average 90 kDa. HSP90 functions primarily as a molecular chaperone, facilitating the folding of other cellular proteins, preventing protein aggregation, or targeting improperly folded proteins to specific degradative pathways. HSP90 is ubiquitously expressed, highly conserved, and accounts for 1-2% of the total cellular protein. Recently introduced, standardized nomenclature has divided the 17 identified HSP90 genes into three related and one unrelated classes, HSP90aa, HSP90AB, HSP90BB, and TRAP, respectively. Six of these genes were functional while the remaining 11 are considered putative pseudogenes. Eukaryotic cells have two principal isoforms of HSP90. The antibody described here was raised against the inducible form, HSP90AA1, a 732 amino acid protein also known as HSP90- α , HSP90A, HSPC1, HSPCA, HSP89- α and LAP2. The constitutively expressed HSP90AB1-isoform 1 is a 724 amino acid protein that is also known as HSP90- β , HSP90B, HSPCB, HSPC2, and HSP89- β . HSP90AB1-1 and HSP90AA share 90% identity. In addition to its role as a molecular chaperone and stress response protein, HSP90 is a central component in a number of basic cellular processes including hormone signaling and cell cycle control.