

DESCRIPTION

Species Reactivity	Human/Mouse
Specificity	Detects human and mouse Leukotriene A4 Hydrolase/LTA4H in direct ELISAs and Western blots.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	<i>S. frugiperda</i> insect ovarian cell line Sf 21-derived recombinant human Leukotriene A4 Hydrolase/LTA4H Pro2-Asp611 Accession # P09960
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Leukotrienes are a family of lipid mediators important in a variety of allergic and inflammatory reactions. Synthesized by leukocytes, these molecules are divided into two classes, the spasmogenic cysteinyl leukotrienes and LTB₄, a classical chemoattractant (1). Encoded by the LTA4H gene, Leukotriene A4 Hydrolase catalyzes the conversion of unstable epoxide LTA₄ to LTB₄, which is the final and committed step in LTB₄ biosynthesis. As a bifunctional zinc metalloenzyme, LTA4H also acts as an arginyl aminopeptidase (2). LTA4H is a drug target for anti-inflammation, and for cancer prevention and therapy (1, 3).

The mature chain of human LTA4H consists of 610 amino acids (residues 2-611). It is highly specific for LTA₄, which also covalently modifies and inhibits the enzyme. The aminopeptidase activity is enhanced by monovalent anions (1). R&D Systems rhLTA4H corresponds to the mature chain, and is characterized by its arginyl aminopeptidase activity.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.