

DESCRIPTION


Species Reactivity	Human/Mouse
Specificity	Detects endogenous human and mouse HSP40 in Western blots.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human HSP40 Met1-Ile340 Accession # P25685
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	1 µg/mL	See Below

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Heat shock protein 40 kDa (HSP40) is the human homologue of the bacterial DnaJ heat shock protein. HSP40, also known as HSPF1 and DnaJ Homolog subfamily B member 1 (DNAJB1), is a 340 amino acid, 40 kDa member of the heat shock protein family. Heat shock proteins (HSPs) are a highly conserved family of stress response proteins. HSPs function primarily as molecular chaperones, facilitating the folding of other cellular proteins, preventing protein aggregation, or targeting improperly folded proteins to specific degradative pathways. Heat Shock Proteins are ubiquitously expressed in all organisms. They are induced in response to various types of environmental stresses like heat, cold, and oxygen deprivation. HSP40 is a stress inducible chaperone that co-localizes with HSP70 and can bind unfolded proteins and prevent protein denaturation and aggregation. The conserved amino terminal J domain can interact with HSP70 and stimulate its ATPase activity. Human HSP40 shares 95% amino acid identity with mouse and rat HSP40.