

DESCRIPTION


Species Reactivity	Human/Mouse
Specificity	Detects endogenous human and mouse Annexin A13 in Western blots. In Western blots, this antibody did not cross-react with recombinant human Annexin A1, A2, A3, A4, A5, A6, A7, A8, A9, A10, or A11.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human Annexin A13 Met1-His316 Accession # P27216
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	1 µg/mL	See Below

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Annexin A13 (ANXA13), also known as Annexin XIII and Intestine-Specific Annexin, is a member of the Annexin protein family. Two isoforms have been identified resulting from alternate splicing. Isoform A consists of 316 amino acids and has a predicted molecular weight of 36 kDa whereas isoform B is composed of 357 amino acids and has a predicted molecular weight of 40 kDa. Annexins are a family of Calcium-dependent phospholipid-binding proteins that are preferentially located on the cytosolic face of the plasma membrane. The Annexin's have a molecular weight of approximately 35 to 40 kDa and consist of a unique amino terminal domain followed by a homologous C-terminal core domain containing the calcium-dependent phospholipid-binding sites. The C terminal domain is comprised of four 60-70 amino acid repeats, known as annexin repeats or an endonexin fold (Annexin A6 contains 8 annexin repeats). The four annexin repeats form a highly α -helical, tightly packed disc known as the annexin domain, which binds to phospholipids in the membrane in a calcium-dependent manner. Members of the annexin family play a role in cytoskeletal interactions, phospholipase inhibition, regulation of cellular growth, and intracellular signal transduction pathways. The specific function of Annexin A13 has not been determined but it is associated with the plasma membrane of undifferentiated, proliferating endothelial cells and differentiated villus enterocytes.