

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Collagen XXIII α 1 in direct ELISAs and Western blots. In direct ELISAs and Western blots, less than 5% cross-reactivity with recombinant human COL25A1 is observed.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant human Collagen XXIII α 1 Glu111-Lys540 Accession # Q86Y22
Conjugate	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Collagen XXIII alpha 1 (sometimes abbreviated COL23A1) is a ~75 kDa type II transmembrane nonfibrillar collagen that is a member of the collagenous transmembrane protein superfamily (1, 2). This family also includes collagens XIII, XVII, XXV and non-collagens with triple-helical regions such as ectodysplasin A, class A macrophage scavenger receptors, and MARCO (2). The human Collagen XXIII mRNA encodes a 540 amino acid (aa) protein containing a 34 aa N-terminal cytoplasmic domain, a 21 aa transmembrane (TM) domain and a 485 aa extracellular domain (ECD). The ECD contains a coiled-coil consensus sequence to aid homotrimerization (aa 64-69), a furin cleavage site (aa 105-110), a pair of cysteines thought to form intermolecular disulfides (aa 106 and 108), and three collagen domains (1, 3-5). The C-terminal 20 aa, including cysteines at aa 525 and 537 of Collagen XXIII, is conserved among TM collagen proteins. Proteolytic cleavage, occurring mainly in the golgi, allows the Collagen XXIII ectodomain to be secreted as a soluble trimer of ~60 kDa subunits (1, 5). Cell surface cleavage can also occur but is slow, presumably due to presence of Collagen XXIII in lipid raft membrane domains (5). The protein database includes three variants of 537, 316 and 309 aa with various portions missing or substituted; all appear to lack TM segments (6). The human Collagen XXIII ECD shares 92%, 93%, 85%, 85% and 84% aa identity with mouse, rat, canine, equine and bovine Collagen XXIII, respectively. Collagen XXIII is concentrated at sites of cell contact in epithelia, and is thought to be an adhesion molecule (2, 4). Its up-regulation has been correlated with aggressiveness in transformed cells, particularly in prostate cancer (1, 7).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.