

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Ficolin-1 in direct ELISAs and Western blots. In direct ELISAs and Western blots, approximately 25% cross-reactivity with recombinant human (rh) Ficolin-2 is observed and less than 1% cross-reactivity with rhFicolin-
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant human Ficolin-1 Gln28-Ala326 Accession # O00602
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Human Ficolin-1 (fibrinogen/collagen-like), also called M-ficolin, is a member of the ficolin family of secreted pattern recognition proteins in the lectin complement activation pathway (1-3). Ficolin-1 is expressed by monocytes, neutrophils and type II alveolar epithelial cells (2-5). It is proposed to be a locally-acting lectin released in a regulated manner (4). Ficolin-1 is not found in plasma, but is detected on the surface of circulating monocytes (4-6). The 35 kDa, 326 amino acid (aa) human Ficolin-1 contains a 28 aa signal sequence, an N-terminal collagen domain and a C-terminal fibrinogen-like (FBG) domain that includes a calcium binding site and one potential N-glycosylation site. Both the collagen and FBG domains mediate trimer formation (7). Like Ficolin-2, larger homo-multimers of Ficolin-1 exist and are likely formed by disulfide bonds at the N-terminus. Sizes corresponding to 12 or 18 subunit oligomers are reported (4, 6, 7). The FBG domain of Ficolin-1 binds microbial ligands that contain acetylated compounds (6). Ligands identified include N-acetyl glucosamine, N-acetyl galactosamine and sialyl-N-acetylglucosamine (4-7). Like other ficolins, Ficolin-1 associates with, and activates the MBL-associated serine protease (MASP) complex, which activates the complement pathway by cleaving C4, contributing to the innate immune response (4-6). Mature human Ficolin-1 shares 76%, 61%, 61%, 76% and 81% aa identity with mouse Ficolin-2 (8), and mouse, rat, canine and porcine Ficolin-1, respectively. It shares 84% and 46% aa identity with human Ficolin-2 and Ficolin-3, respectively. The Ficolin-1 gene is polymorphic, showing at least ten single nucleotide polymorphisms in the promoter and coding regions (1).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.