

Human CEACAM-8/CD66b Antibody


Antigen Affinity-purified Polyclonal Sheep IgG Catalog Number: AF4246

DESCRIPTION			
Species Reactivity	Human		
Specificity	Detects human CEACAM-8/CD66b in direct ELISAs and Western blots. In direct ELISAs, less than 10% cross-reactivity with recombinan human (rh) CEACAM-3, rhCEACAM-6, and rhCEACAM-7 is observed.		
Source	Polyclonal Sheep IgG		
Purification	Antigen Affinity-purified		
Immunogen	E. coli-derived recombinant human CEACAM-8/CD66b Gln35-His141 Accession # P31997		
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.		

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

	Recommended Concentration	Sample	
Western Blot	0.1 μg/mL	Recombinant Human CEACAM-8/CD66b	
Flow Cytometry	0.25 μg/10 ⁶ cells	See Below	
CyTOF-ready	Ready to be labeled with conjugation.	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	



Detection of CEACAM-8/CD66b in Human Granulocytes by Flow Cytometry. Human whole blood granulocytes were stained with Sheep Anti-Human CEACAM-8/CD66b Antigen Affinity-purified Polyclonal Antibody (Catalog # AF-4246, filled histogram) or isotype control antibody (Catalog # 5-001-A, open histogram), followed by NorthernLights™ 637-conjugated Anti-Sheep IgG Secondary Antibody (Catalog # NL011).

PREPARATION AND STORAGE

Reconstitution Reconstitute at 0.2 mg/mL in sterile PBS.

Shipping The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

 * Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 $^\circ$ C

- 12 months from date of receipt, -20 to -70 °C as supplied.
- 1 month, 2 to 8 °C under sterile conditions after reconstitution.
- 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

CEACAM-8 (carcinoembryonic antigen-related cell adhesion molecule 8; also CD66b, CD67 and NCA-95) is a 90 kDa member of the CEACAM subfamily of the CEA family of proteins. It is expressed by neutrophils and eosinophils, and serves as a binding partner for CEACAM-6 and Galectin-3. Mature human CEACAM-8 is a 287 amino acid GPI-linked glycoprotein. It contains one V-type and two C2-type Ig-like domains. No definitive rodent CEACAM-8 has been reported.

Rev. 2/6/2018 Page 1 of 1

