

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human IκB-ε in direct ELISAs. In direct ELISAs, approximately 40% cross-reactivity with recombinant mouse IκB-ε is observed and less than 1% cross-reactivity with recombinant human IκB-β is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human IκB-ε Ser141-Asp500 Accession # O00221
Formulation	Lyophilized from a 0.2 μm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 μm filtered solution in PBS.


APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Immunohistochemistry	5-15 μg/mL	See Below

DATA

Immunohistochemistry



IκB-ε in Human Astrocytoma. IκB-ε was detected in immersion fixed paraffin-embedded sections of human astrocytoma using Goat Anti-Human IκB-ε Antigen Affinity-purified Polyclonal Antibody (Catalog # AF4300) at 3 μg/mL overnight at 4 °C. Before incubation with the primary antibody, tissue was subjected to heat-induced epitope retrieval using Antigen Retrieval Reagent-Basic (Catalog # CTS013). Tissue was stained using the Anti-Goat HRP-DAB Cell & Tissue Staining Kit (brown; Catalog # CTS008) and counterstained with hematoxylin (blue). Specific staining was localized to cytoplasm of neurons. View our protocol for [Chromogenic IHC Staining of Paraffin-embedded Tissue Sections](#).

PREPARATION AND STORAGE

Reconstitution	Sterile PBS to a final concentration of 0.2 mg/mL.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

NFκB inhibitor epsilon (IκB-ε) inhibits the NFκB complex by binding and sequestering it in the cytoplasm. Upon stimulation, IκB-ε is phosphorylated on serine residues marking it for degradation by the ubiquitin pathway. Following IκB-ε degradation, the NFκB complex is able to translocate to the nucleus and activate transcription.