

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human ADAMTS4 in direct ELISAs and Western blots. In direct ELISAs and Western blots, approximately 10% cross-reactivity with recombinant human (rh) ADAMTS5 is observed and less than 1% cross-reactivity with rhADAMTS13, rhADAMTS-L1.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant human ADAMTS4 Phe213-Cys685 Accession # O75173
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunoprecipitation	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

ADAMTS4 (a disintegrin and metalloproteinase with thrombospondin motifs 4), also known as aggrecanase-1, is a member of the family of secreted zinc proteases with a multi-domain structure (1-3). The protein precursors consist of a signal peptide and the following domains: pro, catalytic, disintegrin-like, TS type 1 motif, cysteine-rich, and spacer. It is the only ADAMTS identified that has one TS type I motif. It is an active protease effectively cleaving α-2-macroglobulin and aggrecan at multiple sites, and is inhibited by TIMP-3 with inhibition constants in subnanomolar range (4-6). It receives great attention due to the elevation in its mRNA level after treatment with Interleukin-1 (7). However, in a mouse model of osteoarthritis, ADAMTS4 knock-out mice did not exhibit any significant protective effect (8). The purified rhADAMTS4 starts at the catalytic domain and ends before the spacer domain. If desired, the aggrecanase activity can be inhibited by 5 mM 1,10-phenanthroline and recombinant human TIMP-3 (R&D Systems, Catalog # 973-TM).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.