

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse ECM-1 in direct ELISAs and Western blots. In direct ELISAs and Western blots, approximately 10% cross-reactivity with recombinant human ECM-1 is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse ECM-1 Ala20-Glu559 Accession # AAI38694
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Extracellular matrix protein-1 (ECM-1, ECM-1a) is an 85 kDa, secreted glycoprotein important in connective tissue organization (1-3). Of identified splice variants, the 559 amino acid (aa) form, ECM-1a is most widely expressed, with highest expression in the placenta, heart, and developing bones (3, 4). ECM-1b (434 aa) is found only in tonsil and skin, where it is associated with suprabasal keratinocytes (3, 5). Mouse ECM-1 contains a 19 aa signal peptide and a 540 aa secreted portion that includes an N-terminal proline-rich, cysteine-free region, two tandem repeat domains, and a C-terminal domain. Mature mouse ECM-1 shares 90% aa identity with rat ECM-1 and 65-69% aa identity with corresponding isoforms of human, equine, bovine and canine ECM-1. There are six repeats of a CC(X₇₋₁₀)C motif (x = any aa) within the tandem repeat and C-terminal domains. These motifs, also found in members of the albumin family, are expected to form two (in ECM-1b) or three (in ECM-1a) "double loop" structures that are involved in ligand binding to extracellular matrix molecules such as fibulin-1, perlecan, laminin 332, and fibronectin (4-7). ECM-1 is over-expressed in many malignant epithelial tumors and has demonstrated angiogenic activity (8, 9). A role in regulating alkaline phosphatase during endochondral bone formation has also been suggested (4). In humans, loss of function within the tandem repeat regions due to mutation is considered causative of thickened and irregular extracellular matrix within connective tissue, called lipoid proteinosis (10). Autoantibodies in the skin disease lichen sclerosis also target these repeats (11). The phenotypes of these diseases support a role for ECM-1 as a "biological glue" in the dermis (1, 6, 7).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.