

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse CXCL15/Lungkine in direct ELISAs and Western blots. In direct ELISAs, less than 1% cross-reactivity with rrCINC-1, rmBLC, rmFractalkine, rmGCP-2, rmKC, rmMIG and rmSDF-1α is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant mouse CXCL15/Lungkine (R&D Systems, Catalog # 442-LK) Gln26-Ala167 Accession # Q9WVL7
Conjugate	Alexa Fluor 405 Excitation Wavelength: 405 nm Emission Wavelength: 421 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Mouse Lungkine/CXCL15 (1), also named WECH (2), is a member of the ELR motif-containing CXC chemokines. The cDNA of mouse Lungkine encodes a protein of 166 amino acids (aa) with a 25 aa predicted signal peptide and a 141 aa mature protein with an extremely long C terminal tail that protrudes beyond the chemokine fold. Mouse Lungkine shares 35% aa sequence identity with human ENA-78 and 31% identity with human IL-8. The gene for mouse Lungkine has been mapped to chromosome 5. By Northern blot and in situ hybridization, Lungkine transcripts are only specifically detected in the adult and fetal lung, and its expression is up-regulated under inflammatory conditions (1). Lungkine protein is secreted into bronchoalveolar space and is involved in lung-specific neutrophils trafficking (1). Studies from Lungkine knock out mice suggests that Lungkine is an important mediator of neutrophil migration from the lung parenchyma into the airspace (3). Lungkine is also chemotactic for bone marrow progenitor cells and modulates hematopoietic cell differentiation (2).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.