

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human β -Defensin 3 in direct ELISAs and Western blots. In direct ELISAs and Western blots, less than 1% cross-reactivity with recombinant human β -Defensin 2 is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human β -Defensin 3 Gly23-Lys67 Accession # P81534
Conjugate	Alexa Fluor 405 Excitation Wavelength: 405 nm Emission Wavelength: 421 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

β -Defensin 3, also known as BD3 and DEFB-3, is a membrane-active cationic peptide that functions in inflammation and innate immune responses. There are at least 30 β -Defensins which are distinguished from α -Defensins by the connectivity pattern of their three intramolecular disulfide bonds (1). The 45 amino acid (aa) mature human BD3 shares 38% and 33% aa sequence identity with mouse and rat BD3, respectively (2, 3). It shares 18%-36% aa sequence identity with other human β -Defensins. BD3 is widely expressed among epithelial tissues, notably by keratinocytes and airway epithelial cells. It is up-regulated in response to proinflammatory cytokines, microbial and viral infections, and at the edges of skin wounds (2, 4-6). BD3 induction in osteoarthritis chondrocytes promotes MMP1 and 13 production and inhibits TIMP1 and 2 expression (7). *In vivo* control of BD3 activity is accomplished in part through cleavage by cathepsins B, L, and S (8). BD3 displays strain specific microbicidal activity toward a broad spectrum of bacteria and yeast (2, 9). BD3 also induces monocyte migration, mast cell activation, and a mast cell-dependent increase in vascular permeability (4, 10). Disruption of the intramolecular disulfide bond pattern in BD3 abrogates its monocyte chemoattractant properties but not its antimicrobial properties (11, 12). BD3 inhibits viral infectivity by interacting directly with HIV-1 plus its coreceptor CXCR4 (5, 13), and with HSV glycoprotein B plus its receptor heparan sulfate (14), and by forming a protective coating on the surface of influenza virus target cells (15).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.