

DESCRIPTION

Species Reactivity	Human/Mouse/Rat
Specificity	Detects human, mouse and rat PKCι/λ/ζ in Western blots.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human PKCι Ile455-Val596 Accession # P41743
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Members of the Protein Kinase C (PKC) family are serine/threonine protein kinases that play a key regulatory role in a number of cellular functions including cell growth and differentiation, hormone secretion, and gene expression. Multiple genes and alternative splicing result in three subfamilies, which differ in their co-factor requirements: conventional PKC isoforms (α, βI, βII, and γ) which require calcium and phosphatidylserine (PS), diacylglycerol (DAG) or phorbol esters for activation; novel isoforms (δ, ε, η, and θ), which are calcium-independent but are still regulated by PS, DAG, or phorbol esters; and atypical isoforms (ι, λ, and ζ), which are calcium-independent and do not require PS, DAG, or phorbol esters for activation. PKCι has 72% overall identity to PKCζ. Atypical PKCs have been shown to serve as a convergent downstream target for the PI 3-kinase and TC10 signaling pathways. Stimulation of atypical PKCs by TNF-α has been shown to be required for NF-κB activation. Furthermore, insulin-stimulated atypical PKC activation has been directly implicated in the translocation of GLUT4 and glucose uptake in adipocytes.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.