

**DESCRIPTION**


<b>Species Reactivity</b>	Mouse
<b>Specificity</b>	Detects mouse FPRP in direct ELISAs and Western blots.
<b>Source</b>	Polyclonal Sheep IgG
<b>Purification</b>	Antigen Affinity-purified
<b>Immunogen</b>	Mouse myeloma cell line NS0-derived recombinant mouse FPRP Arg22-Pro832 Accession # NP_035327
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

**APPLICATIONS**

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Western Blot</b>	0.1 µg/mL	Recombinant Mouse FPRP
<b>Flow Cytometry</b>	2.5 µg/10 <sup>6</sup> cells	See Below
<b>CyTOF-ready</b>	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	

**DATA**



**PREPARATION AND STORAGE**

<b>Reconstitution</b>	Reconstitute at 0.2 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>● 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>● 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>● 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

**BACKGROUND**

FPRP (F<sub>2α</sub> Prostaglandin receptor Regulatory Protein; also CD315, EWI-F and CD9 partner-1/P-1) is a 130 kDa type I transmembrane (TM) glycoprotein member of the EWI subfamily, Ig superfamily of molecules. It associates with the 7-TM prostaglandin F<sub>2α</sub> receptor, inhibiting its activity. It also forms complexes with 4-TM CD9 and CD81, linking these molecules to the intracellular actin cytoskeleton. Mature mouse FPRP is 858 amino acids (aa) in length. Its extracellular domain (ECD) contains six C2-type Ig-like domains (aa 22-813) plus a Pro-Thr-Glu-Leu ER retention signal. The ECD of mouse FPRP (aa 22-832) shares 89% and 98% aa identity with the ECD of human and rat FPRP, respectively.