

DESCRIPTION


Species Reactivity	Mouse
Specificity	Detects mouse FPRP in direct ELISAs and Western blots.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse FPRP Arg22-Pro832 Accession # NP_035327
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	0.1 µg/mL	Recombinant Mouse FPRP
Flow Cytometry	2.5 µg/10 ⁶ cells	See Below
CyTOF-ready	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

FPRP (F_{2α} Prostaglandin receptor Regulatory Protein; also CD315, EWI-F and CD9 partner-1/P-1) is a 130 kDa type I transmembrane (TM) glycoprotein member of the EWI subfamily, Ig superfamily of molecules. It associates with the 7-TM prostaglandin F_{2α} receptor, inhibiting its activity. It also forms complexes with 4-TM CD9 and CD81, linking these molecules to the intracellular actin cytoskeleton. Mature mouse FPRP is 858 amino acids (aa) in length. Its extracellular domain (ECD) contains six C2-type Ig-like domains (aa 22-813) plus a Pro-Thr-Glu-Leu ER retention signal. The ECD of mouse FPRP (aa 22-832) shares 89% and 98% aa identity with the ECD of human and rat FPRP, respectively.