

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse Collagen XIII α 1 in direct ELISAs and Western blots. In direct ELISAs, approximately 35% cross-reactivity with recombinant human (rh) COL13A1v4 is observed, and less than 1% cross-reactivity with rhCOL4A1, rhCOL25A1, rhCOL3A1, and rhCOL1A1 i
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse Collagen XIII α 1 Glu107-Gln565 Accession # AAH34164
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Collagen XIII α 1 is an 85-95 kDa protein in the type 2 transmembrane collagen family (1). Mature mouse Collagen XIII α 1 consists of a 40 amino acid (aa) cytoplasmic domain, a 19 aa transmembrane segment, and a 692 aa extracellular domain (ECD). The ECD contains three collagenous regions separated by shorter non-collagenous regions (2, 3). Within comparable regions of the ECD, mouse Collagen XIII α 1 shares 85% and 88% aa sequence identity with human and rat Collagen XIII α 1, respectively. Mouse Collagen XIII α 1 is extensively spliced, with some isoforms showing a tissue specific distribution (2, 4). Collagen XIII α 1 is widely expressed during development and in the adult (4, 5). It localizes to intercellular adherens junctions and cell-matrix focal adhesions (6, 7). Collagen XIII α 1 assembles into disulfide-linked trimers, a process that is enhanced by proline hydroxylation (2, 8). Trimerization involves triple helix formation within the collagenous domains, although portions of the non-collagenous regions can also form coiled coils (8-10). The ECD of trimeric Collagen XIII α 1 is an extended rod-like structure with two flexible hinges that correspond to non-collagenous regions (11). Collagen XIII α 1 clusters in cholesterol-rich domains on the plasma membrane (2, 12), and it can be cleaved from the cell surface or intracellularly by a furin-like protease (12). Collagen XIII α 1 binds the extracellular matrix molecules fibronectin, heparin, integrin α 1, nidogen-2, and perlecan (11, 13). The shed ECD retains its ability to bind fibronectin and can interfere with matrix formation (14).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.