

DESCRIPTION

Species Reactivity	Human
Specificity	Detects endogenous human DFF45 in Western blots.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human DFF45 Met1-Thr331 Accession # O00273
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

DFF45 (DNA fragmentation factor 45 kDa subunit, also known as ICAD) is a 45 kDa member of the CIDE domain-containing family of proteins. Human DFF45 is 331 amino acids (aa) in length. It contains an N-terminal CIDE domain (aa 17-96) plus two caspase-3 cleavage sites (aa's 117-118 and 224-225). DFF45 is a cytoplasmic protein that inhibits DFF40 DNAase. Normally, DFF45 and DFF40 form an enzymatically-inactive noncovalent heterodimer. Upon activation of the apoptotic cascade, DFF45 is cleaved into three fragments. These dissociate from DFF40 and induce DFF40 oligomerization and activation. Human DFF45 has two potential splice forms. One is truncated, and shows a seven aa substitution for aa 262-331; a second is extended, and shows a 45 aa substitution for the C-terminal five amino acids. Human DFF45 is 76% aa identical to mouse DFF45.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.