

## DESCRIPTION

|                                                                                                                                                                                                         |                                                                                                                                                      |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Species Reactivity</b>                                                                                                                                                                               | Mouse                                                                                                                                                |
| <b>Specificity</b>                                                                                                                                                                                      | Detects mouse SCARA5 in direct ELISAs and Western blots. In direct ELISAs, less than 40% cross-reactivity with recombinant human SCARA5 is observed. |
| <b>Source</b>                                                                                                                                                                                           | Polyclonal Sheep IgG                                                                                                                                 |
| <b>Purification</b>                                                                                                                                                                                     | Antigen Affinity-purified                                                                                                                            |
| <b>Immunogen</b>                                                                                                                                                                                        | Chinese hamster ovary cell line CHO-derived recombinant mouse SCARA5<br>Arg83-Pro491<br>Accession # Q8K299                                           |
| <b>Conjugate</b>                                                                                                                                                                                        | Alexa Fluor 488<br>Excitation Wavelength: 488 nm<br>Emission Wavelength: 515-545 nm                                                                  |
| <b>Formulation</b>                                                                                                                                                                                      | Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide                                                                                         |
| *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions. |                                                                                                                                                      |

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

|                       |                                                                        |
|-----------------------|------------------------------------------------------------------------|
| <b>CyTOF-ready</b>    | Optimal dilution of this antibody should be experimentally determined. |
| <b>Western Blot</b>   | Optimal dilution of this antibody should be experimentally determined. |
| <b>Flow Cytometry</b> | Optimal dilution of this antibody should be experimentally determined. |

## PREPARATION AND STORAGE

|                                |                                                                                                                   |
|--------------------------------|-------------------------------------------------------------------------------------------------------------------|
| <b>Shipping</b>                | The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below. |
| <b>Stability &amp; Storage</b> | Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied                          |

## BACKGROUND

Scavenger receptor A5 (SCARA5; also known as testis expressed scavenger receptor Tesr) is a Class A scavenger receptor that is related to CL-P1, MARCO, SCARA3, and SR-A1. All are type II transmembrane proteins that contain a collagenous stalk (1-3). Mature mouse SCARA5 consists of a 60 amino acid (aa) cytoplasmic domain, a 21 aa transmembrane segment, and a 410 aa C-terminal extracellular domain (ECD) that contains a coiled-coil, a collagen-like, and a scavenger receptor cysteine-rich (SRCR) region (4, 5). Within the ECD, mouse SCARA5 shares 87% and 98% aa sequence identity with human and rat SCARA5, respectively. It shares 24%-29% aa sequence identity with CL-P1, MARCO, SCARA3, and SR-A1. Alternative splicing generates one isoform that lacks the SRCR domain and a second that additionally lacks the cytoplasmic and transmembrane regions. SCARA5 is a cell surface disulfide-linked homotrimer of > 250 kDa (5). It is highly expressed in testicular Sertoli and germ cells and more weakly in the epithelia of other tissues (4, 5). During mouse development, SCARA5 expression is attenuated in female embryos at the time of sex determination, whereas it is maintained in the developing testis (4). SCARA5 is re-expressed in the adult ovary (4). SCARA5 binds heat-killed bacterial particles but not yeast particles or modified LDL (5).

## PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.