

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human CDP/CUTL1 in direct ELISAs and Western blots.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human CDP/CUTL1 Ile64-Ser268 Accession # P39880
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

CDP (CCAAT Displacement Protein; also CUX-1 and CUTL-1) is a 180-190 kDa member of the CUT homeobox family of proteins. It is found in multiple cell types, and appears to act as a negative regulator of transcription for genes as diverse as c-myc, CD8 and TGF-βRII. Repression is assumed to occur either by direct DNA interaction, which interferes with transcriptional activator binding to gene promoters, or through the recruitment of active HDACs to target gene promoter sequences. Human CDP is 1505 amino acids (aa) in length. It contains three consecutive DNA-binding CUT domains (aa 542-629; 934-1007; 1125-1202) plus one C-terminal homeobox domain (aa 1244-1303). CDP undergoes phosphorylation, resulting in its inactivation, and reportedly also homodimerizes. There are multiple splice variants (termed CASP), ranging from 75-85 kDa in size. Most are characterized by the presence of a unique 260 aa substitution for aa 410-1505 that contains a Golgi transmembrane domain. CDP also undergoes extensive proteolytic processing, creating functional DNA binding C-terminal fragments that range from 75-110 kDa in size. Over aa 64-268, human CDP shares 97% aa identity with mouse CDP.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.