

DESCRIPTION

Species Reactivity	Human/Mouse
Specificity	Detects human and mouse Neurturin in direct ELISAs and Western blots.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant mouse Neurturin
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Blockade of Receptor-ligand Interaction	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Neurturin is a member of the GDNF family of ligands, which include glial cell-derived neurotrophic factor (GDNF), Neurturin, Persephin, and Artemin. GDNF family proteins are distant members of the Transforming Growth Factor β (TGF- β) superfamily (1 - 4). Similar to other TGF- β family proteins, Neurturin is synthesized as a precursor protein that is cleaved at the dibasic cleavage site (RXXR) to release the carboxy-terminal domain. The carboxy-terminal domain of Neurturin contains the characteristic seven conserved cysteine residues necessary for the formation of the cysteine knot and the single interchain disulfide bond. Biologically active Neurturin is a disulfide-linked homodimer of the carboxy-terminal 100 amino acid residues. Mature mouse Neurturin shares 91% amino acid sequence similarity with human Neurturin. Mature Neurturin also shares about 40% similarities with the other three members of the GDNF family ligands (2, 5). Unlike other members of TGF- β family, bioactivities of all GDNF family ligands are mediated through a unique multicomponent receptor complex composed of high affinity ligand binding component (GFR α -1 - GFR α -4) and a common signaling component (cRET receptor tyrosine kinase). Each member of the GDNF family ligands has its preferred binding protein. Neurturin preferentially binds to GFR α -2 but can also bind GFR α -1 at higher concentration (5 - 8). Neurturin has been shown to promote the survival of a variety of neurons including sympathetic, sensory, and central nervous system neurons. Neurturin is expressed in both neuronal and non-neuronal tissues. It may play a role in regulating the development and maintenance of the central and peripheral nervous systems as well as non-neuronal systems (9).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.