

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human SPRED2 in direct ELISAs and Western blots. In direct ELISAs, approximately 10% cross-reactivity with recombinant human (rh) SPRED1 is observed and less than 3% cross-reactivity with rhSPRED4 is observed.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human SPRED2 Lys117-Ser297 Accession # Q7Z698
Conjugate	Alexa Fluor 594 Excitation Wavelength: 590 nm Emission Wavelength: 617 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunocytochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

SPRED2 (Sprouty-related protein with an EVH1 domain-2) is a 47-53 kDa member of the SPRED family of regulatory molecules. In human, it is found in keratinocytes, intestinal columnar epithelium, and salivary duct epithelium. It inhibits receptor tyrosine kinase (RTK) signaling by blocking Ras activation of Raf. This is achieved by inducing the lysosomal degradation of RTKs following SPRED2 binding to NBR1. Human SPRED2 is 418 amino acids (aa) in length. It contains an N-terminal WH1/EVH1 domain that is involved in ERK inhibition (aa 5-122), a central KBD domain that binds to the SCFR (aa 201-257), and a C-terminal Sprouty-related domain that mediates homodimerization, and heterodimerization with SPRED1 (aa 308-416). SPRED is ubiquitinated, and undergoes phosphorylation on Ser, Thr and Tyr residues. There is one splice variant that shows a six aa substitution for aa 1-9. Over aa 117-297, human SPRED2 shares 84% aa identity with mouse SPRED2.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.