

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse ASAH1/N-acyl ethanolamine-hydrolyzing Acid A in direct ELISAs and Western blots. In direct ELISAs, approximately 10% cross-reactivity with recombinant human (rh) ASAH1 is observed and less than 5% cross-reactivity rhASAH1-
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse ASAH1/N-acyl ethanolamine-hydrolyzing Acid A Val33-Ser362 Accession # AAH04572
Conjugate	Alexa Fluor 594 Excitation Wavelength: 590 nm Emission Wavelength: 617 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunoprecipitation	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

The mouse ASAH1 gene encodes N-acyl ethanolamine-hydrolyzing Acid Amidase (NAAA), a fatty acid amidase with maximal activity at acidic pH (1). NAAA hydrolyzes a number of N-acyl ethanolamines, including N-myristoyl-, N-stearoyl-, N-oleoyl-, and N-arachidonoyl, but is most active against N-palmitoylethanolamine (2). NAAA is a member of the cholesteryl glycerol hydrolase family of enzymes, and is structurally similar to acid ceramidase (1). NAAA is both a lysosomal and a secreted enzyme, and like acid ceramidase, has been observed to be proteolytically processed during maturation (1). Through its amidase activity, ASAH1 may play a role in the termination of the actions of a variety of N-acyl ethanolamides (3). NAAA can be distinguished from anandamide amidohydrolase by its lack of inhibition by methyl arachidonoyl fluorophosphonate (2).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.