

DESCRIPTION

Species Reactivity	Human/Mouse
Specificity	Detects human and mouse LRRTM3 in direct ELISAs and Western blots. In direct ELISA, approximately 25% cross-reactivity with recombinant human (rh) LRRTM4, and less than 5% cross-reactivity with rhLRRTM1 and rhLRRTM2 is observed.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant human LRRTM3 Glu31-Lys419 Accession # Q86VH5
Conjugate	Alexa Fluor 350 Excitation Wavelength: 346 nm Emission Wavelength: 442 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Human LRRTM3 (leucine-rich repeat transmembrane neuronal 3) is a 63 kDa (predicted) type I transmembrane protein, and one of four members of the LRRTM family of proteins within the leucine-rich repeat (LRR) superfamily (1). There are two isoforms of LRRTM3. Isoform 1 is synthesized as a precursor with a 30 amino acid (aa) signal sequence, a 389 aa extracellular region, a 21 aa transmembrane region, and a 141 aa cytoplasmic region. In isoform 2, aa 513 is changed from isoleucine to valine, and aa 514 to 581 are missing, producing a cytoplasmic region of only 74 aa. The extracellular region of both isoforms contains one N-linked glycosylation site, a leucine-rich repeat N-terminal domain bordered by four conserved cysteines, and 10 LRRs flanked by cysteine-rich domains (1). The cytoplasmic region of both isoforms contains several tyrosine, serine, and threonine residues that have the potential to be phosphorylated, and thus to be involved in signal transduction (1). The C-terminal of isoform 2 also contains a conserved glutamic acid-cysteine-glutamic acid-valine sequence for potential interaction with PDZ proteins (1, 2). Mature human LRRTM3 (isoform 1) is 98% identical to mouse LRRTM3. In the mouse, beginning at 8.5 dpc, strong levels of LRRTM3 can be detected in the neural progenitors of the neural plate that will develop into the rostral neural tube, the forebrain, a stripe in the hindbrain, and the region of the presomitic mesoderm/somite boundary (2). By 15 dpc, LRRTM3 is expressed broadly and accumulates thereafter (3). In the adult, LRRTM3 is expressed almost exclusively in the brain with high expression in the cortical laminae and dentate gyrus, as well as detectable levels in the hypothalamus and amygdala (3). Functionally, LRRTM3 may be involved in the formation of the CNS and maintenance of CNS structure and function in the adult brain (1). In addition, LRRTM3 has been shown to promote processing of amyloid-precursor protein by BACE1, and is a positional candidate gene for late-onset Alzheimer's disease (3).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.