

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human ITM2C in direct ELISAs and Western blots. In direct ELISAs and Western blots, less than 1% cross-reactivity with recombinant human (rh) ITM2A and rhITM2B is observed.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human ITM2C Ala86-Val267 Accession # Q9NQX7
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

ITM2C (Integral membrane protein 2C; also BRI3 and cerebral protein 14) is a 36-38 kDa member of the BRI₃ family of proteins. It is found on neurons and appears to interact with APP, where ITM2C binding blocks proteolytic processing of APP, plus SCG10, where ITM2C binding blocks the microtubule-destabilizing activity of SCG10, thus inhibiting neurite outgrowth. Human ITM2C is a 267 amino acid (aa) type II transmembrane glycoprotein. Proteolytic cleavage by furin at Arg242Gly243 may create a truncated 34-35 kDa ITM2C plus a C-terminal 4 kDa active peptide (aa 243-267). The ECD (aa 76-267) contains one BRICHOS domain (aa 136-230). ITM2C may exist as a dimer. There are two splice variants, one that shows a deletion of aa 41-87, and a second that shows a deletion of aa 151-187. Over aa 86-267, human ITM2C shares 96% aa identity with mouse ITM2C.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.