

DESCRIPTION


Species Reactivity	Human
Specificity	Detects human ERMAP in direct ELISAs and Western blots.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant human ERMAP His30-Ala155 Accession # Q96PL5
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	0.1 µg/mL	Recombinant Human ERMAP
Flow Cytometry	2.5 µg/10 ⁶ cells	See Below
CyTOF-ready	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

ERMAP (erythrocyte membrane-associated protein; also Scianna blood group antigen) is a 60-66 kDa member of the BTN/MOG family, Ig-Superfamily of proteins. It is expressed on erythrocytes and erythrocyte precursors and likely serves as a cell adhesion molecule. Mature human ERMAP is a 446 amino acid (aa) type I transmembrane glycoprotein. It possesses a 126 aa extracellular domain (ECD) (aa 30-155) that contains one V-type Ig-like domain (aa 30-140). Single aa changes at Gly35, Glu47, Gly57, Pro60 and Arg81 generate distinct antigens of the Scianna blood group. There is one alternate start site at Met91 and a splice variant that shows an 11 aa substitution for aa 103-475. Over aa 30-155, human ERMAP is 56% aa identical to mouse ERMAP. In contrast to human ERMAP ECD, mouse REMAP ECD contains one extra Ig-like domain.