

DESCRIPTION



Species Reactivity	Human/Mouse/Rat
Specificity	Detects human, mouse, and rat NEDD8 in direct ELISAs and Western blots. In direct ELISAs and Western blots, less than 2% cross-reactivity with recombinant human NEDD9 is observed.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human NEDD8 Leu2-Gly77 Accession # Q15843
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	2 µg/mL	See Below
Immunohistochemistry	5-15 µg/mL	See Below

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> • 12 months from date of receipt, -20 to -70 °C as supplied. • 1 month, 2 to 8 °C under sterile conditions after reconstitution. • 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

NEDD8 (Neural precursor cell-expressed developmentally down-regulated gene 8; also Rub1) is a 6-8 kDa member of the ubiquitin family of proteins. It is expressed in striated muscle, and via its C-terminal Gly, forms covalent bonds with cullin family proteins. This conjugation generates ubiquitin ligase activity that regulates cell cycle proteins. Human pro-NEDD8 is 81 amino acids (aa) in length. It contains one ubiquitin-like domain (aa 1-65), and a 4-5 aa C-terminal propeptide that is cleaved to expose a C-terminal Gly residue that is used to generate a Gly-Lys intermolecular bond. Mature human NEDD8 shows 100% aa sequence identity to NEDD8 from both mouse and canine.