Human Ikaros Alexa Fluor® Plus 405-conjugated Antibody

Antigen Affinity-purified Polyclonal Goat IgG
Catalog Number: AF4984AFP405
100 µg

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Ikaros in direct ELISAs and Western blots. In direct ELISAs and Western blots, less than 1% cross-reactivity with recombinat human (rh) ZIC-1, rhZNF-24, and rhZNF-206 is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	*E. coli*-derived recombinant human Ikaros Ser427-Ser519 Accession # Q13422
Conjugate	Alexa Fluor Plus 405 Excitation Wavelength: 404 nm Emission Wavelength: 455 nm
Formulation	Supplied 0.2 mg/mL in a saline solution containing BSA and Sodium Azide. *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: *Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.*

CyTOF-ready	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.
Chromatin Immunoprecipitation (ChIP)	Optimal dilution of this antibody should be experimentally determined.
Immunocytochemistry	Optimal dilution of this antibody should be experimentally determined.
Intracellular Staining by Flow Cytometry	Optimal dilution of this antibody should be experimentally determined.

DATA

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Ikaros (also LyF1) is a 60 kDa member of the C2H2-type zinc-finger protein family. It is found in both T and B cells and serves as a context-dependent activator or repressor of genes. Human Ikaros is 519 amino acids (aa) in length. It possesses two zinc-finger domains, one at the N-terminus that contains four zinc-finger motifs (aa 117-224) and one at the C-terminus that contains two zinc-finger motifs (aa 462-514). The C-terminal motifs mediate homo- or heterodimerization, while the N-terminal motifs bind DNA. Multiple splice forms of 37 kDa to 47 kDa exist (Ikaros 2-8) with reduced numbers of N-terminal zinc-finger motifs. At least three are needed for DNA binding, and a heterodimer with a two-motif monomer is suggested to be transcriptionally inert. Over aa 427-519 (which are not spliced), human Ikaros shows 88% aa identity to mouse Ikaros.

PRODUCT SPECIFIC NOTICES

This product is provided under an intellectual property license from Life Technologies Corporation. The transfer of this product is conditioned on the buyer using the purchased product solely in research conducted by the buyer, excluding contract research or any fee for service research, and the buyer must not (1) use this product or its components for (a) diagnostic, therapeutic or prophylactic purposes; (b) testing, analysis or screening services, or information in return for compensation on a per-test basis; or (c) manufacturing or quality assurance or quality control, and/or (2) sell or transfer this product or its components for resale, whether or not resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5781 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@thermofisher.com.

Bio-Techne®
Global | bio-techne.com info@bio-techne.com techsupport@bio-techne.com TEL: 1.612.379.2956
USA | TEL: 800.343.7475 Canada | TEL: 855.668.8722 Europe | Middle East | Africa TEL: +44.0.1235.529449
China | info.cn@bio-techne.com TEL: 400.821.3475