

## DESCRIPTION

|                           |                                                                                                                                                                                                                                                                                                        |
|---------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Species Reactivity</b> | Rat                                                                                                                                                                                                                                                                                                    |
| <b>Specificity</b>        | Detects rat IL-1 $\alpha$ /IL-1F1 in direct ELISAs and Western blots. In direct ELISAs, less than 50% cross-reactivity with recombinant mouse IL-1 $\alpha$ /IL-1F1 and recombinant cotton rat IL-1 $\alpha$ /IL-1F1 is observed, and less than 15% cross-reactivity with human IL-1 $\alpha$ /IL-1F1. |
| <b>Source</b>             | Polyclonal Goat IgG                                                                                                                                                                                                                                                                                    |
| <b>Purification</b>       | Antigen Affinity-purified                                                                                                                                                                                                                                                                              |
| <b>Immunogen</b>          | <i>E. coli</i> -derived recombinant rat IL-1 $\alpha$ /IL-1F1<br>Ser115-Ser270<br>Accession # P16598                                                                                                                                                                                                   |
| <b>Conjugate</b>          | Alexa Fluor 750<br>Excitation Wavelength: 749 nm<br>Emission Wavelength: 775 nm                                                                                                                                                                                                                        |
| <b>Formulation</b>        | Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide<br><br>*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.                            |

## APPLICATIONS

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

|                       |                                                                        |
|-----------------------|------------------------------------------------------------------------|
| <b>Neutralization</b> | Optimal dilution of this antibody should be experimentally determined. |
| <b>Western Blot</b>   | Optimal dilution of this antibody should be experimentally determined. |

## PREPARATION AND STORAGE

|                                |                                                                                                                   |
|--------------------------------|-------------------------------------------------------------------------------------------------------------------|
| <b>Shipping</b>                | The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below. |
| <b>Stability &amp; Storage</b> | Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied                          |

## BACKGROUND

Interleukin 1 (IL-1) is a name that designates two proteins, IL-1 $\alpha$  and IL-1 $\beta$ , which are the products of distinct genes, but which show approximately 25% amino acid (aa) sequence identity and which recognize the same cell surface receptors. Although IL-1 production is generally considered to be a consequence of inflammation, evidence suggests that IL-1 is also temporally upregulated during bone formation and the menstrual cycle and can be induced in response to nervous system stimulation. In response to classic stimuli produced by inflammatory agents, infections or microbial endotoxins, a dramatic increase in the production of IL-1 by macrophages and various other cells is seen. Cells in particular known to produce IL-1 include osteoblasts, monocytes, macrophages, keratinocytes, Kupffer cells, hepatocytes, thymic and salivary gland epithelium, Schwann cells, fibroblasts, and glia (oligodendroglia, astrocytes, and microglia).

IL-1 $\alpha$  and IL-1 $\beta$  are both synthesized as 31 kDa precursors that are subsequently cleaved into proteins with molecular weights of approximately 17,000 Da. Neither precursor contains a typical hydrophobic signal peptide sequence and most of the precursor form of IL-1 $\alpha$  remains in the cytosol of cells, although there is evidence for a membrane-bound form of the precursor form of IL-1 $\alpha$ . The IL-1 $\alpha$  precursor reportedly shows full biological activity in the EL-4 assay. Among various species, the aa sequence of mature IL-1 $\alpha$  is conserved 60-70% and human IL-1 has been found to be biologically active on murine cell lines. Both forms of IL-1 bind to the same receptors, designated type I and type II. Evidence suggests that only the type I receptor is capable of signal transduction and that the type II receptor may function as a decoy, binding IL-1 and thus preventing binding of IL-1 to the type I receptor.

## PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.