

**DESCRIPTION**

<b>Species Reactivity</b>	Mouse/Rat
<b>Specificity</b>	Detects mouse and rat PRAT4B in Western blots. In direct ELISAs, less than 5% cross-reactivity with recombinant human PRAT4B and recombinant mouse PRAT4A is observed.
<b>Source</b>	Polyclonal Goat IgG
<b>Purification</b>	Antigen Affinity-purified
<b>Immunogen</b>	Mouse myeloma cell line NS0-derive recombinant mouse PRAT4B Thr30-Leu245 Accession # Q8BQ47
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.


**APPLICATIONS**

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Western Blot</b>	1 µg/mL	See Below
<b>Flow Cytometry</b>	2.5 µg/10 <sup>6</sup> cells	See Below
<b>CyTOF-ready</b>	Ready to be labeled using established conjugation methods. No BSA or other carrier proteins that could interfere with conjugation.	


**DATA**

**Western Blot**



**Detection of Mouse and Rat PRAT4B by Western Blot.**  
Western blot shows lysates of rat lung tissue and mouse spleen tissue. PVDF Membrane was probed with 1 µg/mL of Goat Anti-Mouse/Rat PRAT4B Antigen Affinity-purified Polyclonal Antibody (Catalog # AF5015) followed by HRP-conjugated Anti-Goat IgG Secondary Antibody (Catalog # HAF019). A specific band was detected for PRAT4B at approximately 32 kDa (as indicated). This experiment was conducted under reducing conditions and using Immunoblot Buffer Group 1.

**Flow Cytometry**



**Detection of PRAT4B in RAW264.7 Mouse Cell Line by Flow Cytometry**  
RAW 264.7 mouse monocyte/macrophage cell line was stained with Goat Anti-Mouse/Rat PRAT4B Antigen Affinity-purified Polyclonal Antibody (Catalog # AF5015, filled histogram) or control antibody (Catalog # AB-108-C, open histogram), followed by Allophycocyanin-conjugated Anti-Goat IgG Secondary Antibody (Catalog # F0108).

**PREPARATION AND STORAGE**

<b>Reconstitution</b>	Reconstitute at 0.2 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>• 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>• 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>• 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

**BACKGROUND**

PRAT4B (protein associated with TLR4-B; also Cnpy4, protein canopy homolog 4) is a member of the canopy family of proteins. It is expressed by diverse cell types, including macrophages, fibroblasts and B cells. PRAT4B appears to be both secreted, and retained internally, where it may stabilize immature TLR protein expression in the ER, and impact its appearance on the cell surface. Mouse PRAT4B is 245 amino acids (aa) in length. It contains a 27 aa signal sequence plus one coiled-coil region (aa 209 - 237). There is one potential splice variant that shows a 13 aa substitution for the C-terminal nine amino acids. Over aa 30 - 245, mouse PRAT4B is 95% and 87% aa identical to rat and human PRAT4B, respectively.