

**DESCRIPTION**


<b>Species Reactivity</b>	Human/Mouse
<b>Specificity</b>	Detects endogenous human and mouse TIAM1 in Western blots.
<b>Source</b>	Polyclonal Sheep IgG
<b>Purification</b>	Antigen Affinity-purified
<b>Immunogen</b>	<i>E. coli</i> -derived recombinant human TIAM1 Met1-Ala378 Accession # Q13009
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

**APPLICATIONS**

**Please Note:** Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	<b>Recommended Concentration</b>	<b>Sample</b>
<b>Western Blot</b>	1 µg/mL	See Below

**DATA**



**PREPARATION AND STORAGE**

<b>Reconstitution</b>	Reconstitute at 0.2 mg/mL in sterile PBS.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
<b>Stability &amp; Storage</b>	<b>Use a manual defrost freezer and avoid repeated freeze-thaw cycles.</b> <ul style="list-style-type: none"> <li>• 12 months from date of receipt, -20 to -70 °C as supplied.</li> <li>• 1 month, 2 to 8 °C under sterile conditions after reconstitution.</li> <li>• 6 months, -20 to -70 °C under sterile conditions after reconstitution.</li> </ul>

**BACKGROUND**

TIAM1 (T lymphoma invasion and metastasis) is an approximately 200 kDa member of the TIAM family of guanine-nucleotide exchange factors. It is expressed in multiple cell types, including endothelial cells, neurons, Schwann cells and almost all tumors. TIAM1 stimulates the activity of RHO-like GTPases and thus connects extracellular signals to the cytoskeleton. Human TIAM1 is 1591 amino acids (aa) in length. It contains a myristoylation site at Gly2, plus two pleckstrin homology domains (aa 434 - 549 and 1261 - 1397), a Ras-binding region (aa 765 - 832) and a RhoGEF/DH domain (aa 1041 - 1232). Over aa 1 - 378, human TIAM1 shows 94% aa identity to mouse TIAM1.