

DESCRIPTION


Species Reactivity	Human/Mouse
Specificity	Detects human and mouse Bcl-6 in Western blots.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant mouse Bcl-6 Glu522-Thr696 Accession # Q8CB25
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. *General Protocols* are available in the *Technical Information* section on our website.

	Recommended Concentration	Sample
Western Blot	0.1 µg/mL	See Below

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> ● 12 months from date of receipt, -20 to -70 °C as supplied. ● 1 month, 2 to 8 °C under sterile conditions after reconstitution. ● 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

Bcl-6 (B-cell lymphoma protein 6; also ZFP51, LAZ-3 and Bcl-5) is an 80 kDa member of the Kruppel-like family of zinc finger proteins. It is expressed in germinal center B cells, neutrophils, monocytes, and immature dendritic cells, and serves as a potent transcriptional repressor of genes such as p53. Human Bcl-6 is 706 amino acids (aa) in length and contains an N-terminal BTB/POZ protein interaction domain (aa 32-99) and a series of six C2H2-type C-terminal zinc finger domains that bind DNA (aa 518-681). Centrally located are three PEST sequences (aa 336-430) that undergo phosphorylation by MAPK. This increases the molecular weight of Bcl-6 to 97 kDa and induces its degradation. Human Bcl-6 is about 99% aa identical to mouse and canine Bcl-6.