

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human ADAM19 in direct ELISAs and Western blots. In direct ELISAs and Western blots, approximately 25% cross-reactivity with recombinant mouse ADAM19 is observed, and less than 5% cross-reactivity with recombinant human (rh)&nb
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant human ADAM19 Arg26-Gln501 (Ser32 del) (Lys33Arg) Accession # Q9H013
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot	Optimal dilution of this antibody should be experimentally determined.
Immunoprecipitation	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

ADAM19 (a disintegrin and metalloprotease 19; also MADDAM and Meltrin-β) is a 95-100 kDa member of the M12B peptidase family of enzymes. It is expressed on multiple cell types including monocytes, fibroblasts, osteoblasts and dendritic cells. The human ADAM19 proenzyme is a 931 amino acid (aa) type I transmembrane protein. It contains a 178 aa cleavable proregion (aa 26-203) and a 497 aa extracellular domain (ECD) (aa 204-702). The ECD contains a metalloprotease domain (aa 211-409), a disintegrin region (aa 417-503) and a Cys-rich domain (aa 504-650). The ECD circulates in plasma bound to 700 kDa α2-Macroglobulin. One splice variant shows an alternate start site at Met71, while two isoforms show variations after Lys902. Over aa 205-702, human ADAM19 is 87% aa identical to mouse ADAM19.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.