

## Human Protocadherin α1 Alexa Fluor® 594-conjugated Antibody

Antigen Affinity-purified Polyclonal Sheep IgG Catalog Number: AF5064T

100 µg

DESCRIPTION		
Species Reactivity	Human	
Specificity	Detects human Protocadherin alpha 1 (PCDHα1) in direct ELISAs and Western blots.	
Source	Polyclonal Sheep IgG	
Purification	Antigen Affinity-purified	
Immunogen	S. frugiperda insect ovarian cell line $Sf$ 21-derived recombinant human Protocadherin $\alpha$ 1 Gln30-Pro689 Accession # Q9Y513	
Conjugate	Alexa Fluor 594 Excitation Wavelength: 590 nm Emission Wavelength: 617 nm	
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide	
	*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS			
Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.			
Western Blot	Optimal dilution of this antibody should be experimentally determined.		
Immunohistochemistry	Optimal dilution of this antibody should be experimentally determined.		

PREPARATION AND STORAGE		
Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.	
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied	

## **BACKGROUND**

Protocadherin α1 (PCDHα1) is a group A member of the Protocadherin family of molecules. It is expressed in synaptic membranes of neurons, forms obligate Ca<sup>++</sup> independent complexes with various PCDH γ family members and demonstrates heterophilic binding activity towards β1 integrins. Human PCDHα1 is 921 amino acids (aa) in length. It is a type I transmembrane glycoprotein that contains a 668 aa extracellular domain (ECD) plus a 232 aa cytoplasmic region. Both regions contribute to PCDH-γ binding. There are six cadherin domains in the ECD. ADAM10 cleavage generates an 80-90 kDa soluble fragment and a soluble splice form exists that is missing aa 535-798. Over aa 30-689, human PCDHα1 shares 83% aa identity with mouse PCDHα1.

## PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.

Rev. 9/16/2025 Page 1 of 1