

Human IQGAP1 Antibody

Antigen Affinity-purified Polyclonal Sheep IgG Catalog Number: AF5069


DESCRIPTION	
Species Reactivity	Human
Specificity	Detects endogenous human IQGAP1 in Western blots.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	E. coli-derived recombinant human IQGAP1 Lys505-Ser717 Accession # P46940
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Western Blot	1 μg/mL	See Below

DATA



Detection of Human IQGAP1 by Western Blot.

Western blot shows lysates of HUVEC human umbilical vein endothelial cells, HepG2 human hepatocellular carcinoma cell line, U937 human histiocytic lymphoma cell line, and MRC-5 human embryonic lung fibroblast cell line. PVDF membrane was probed with 1 μ g/mL of Human IQGAP1 Antigen Affinity-purified Polyclonal Antibody (Catalog # AF5069) followed by HRP-conjugated AntiSheep IgG Secondary Antibody (Catalog # HAF016). A specific band was detected for IQGAP1 at approximately 195 kDa (as indicated). This experiment was conducted under reducing conditions and using Immunoblot Buffer Group 1.

PKE	PAKA	HON	AND	SIC	JKAC	jΕ

Reconstitution Reconstitute at 0.2 mg/mL in sterile PBS.

Shipping The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below.

*Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C

Stability & Storage Use a manual defrost freezer and avoid repeated freeze-thaw cycles.

- 12 months from date of receipt, -20 to -70 °C as supplied.
- 1 month, 2 to 8 °C under sterile conditions after reconstitution
- 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

IQGAP1 (IQ motif with Ras <u>GAP</u>-related domain <u>1</u>) is a 195 kDa member of the IQGAP family of proteins. It serves as an effector protein for Rho family members, and regulates actin polymerization. Human IQGAP1 is 1657 amino acids (aa) in length. It contains one actin-binding CH domain (aa 45-159), a Pro-binding WWP region (aa 685-711), four consecutive calmodulin-binding IQ motifs (aa 745-864), and a noncatalytic Ras-GAP domain (aa 1004-1237). Amino acids 763-863 mediate homodimerization. There are at least three IQGAP1 splice variants. One shows a two Lys substitution for aa 270-1657, a second shows a ten Lys substitution for aa 299-1657, and a third shows a three Lys substitution for aa 941-1657.

Rev. 2/6/2018 Page 1 of 1

