

DESCRIPTION

Species Reactivity	Human/Mouse/Rat
Specificity	Detects human, mouse, and rat LMW-PTP/ACP1 in Western blots.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human LMW-PTP/ACP1 Ala2-His158 Accession # P24666
Conjugate	Alexa Fluor 647 Excitation Wavelength: 650 nm Emission Wavelength: 668 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

LMW-PTP (Low Molecular Weight Protein Tyrosine Phosphatase; also Red cell acid phosphatase 1/ACP1, HCPTP, and HAAP/Adipocyte acid phosphatase) is a 17-18 kDa member of the LMWPTP family, PTP superfamily of enzymes. It is widely expressed, being found in cells diverse as B and T cells, endothelial cells and vascular smooth muscle cells. LMW-PTP dephosphorylates multiple substrates, often in a cell-dependent manner. Among the substrates for LMW-PTP include PDGRβ, EphA2 and β-catenin. PMW-PTP is found intracellularly, generally associated with two cytoplasmic pools, one in the cytosol and the other accompanying the cytoskeleton. Human LMW-PTP is 158 amino acids (aa) in length. The catalytic region spans aa 9-154, and there are two acetylation sites at Ala2 and Lys156. Phosphorylation on Tyr132 and Tyr133 occurs on cytoskeleton-associated LMW-PTP, and has the effect of increasing its phosphatase activity. There are three alternative splice variants reported for LMW-PTP. One shows a 34 aa substitution for aa 41-74, a second shows a deletion of aa 41-74, while a third contains a 35 aa substitution for aa 78-158. The last two isoforms may function as negative regulators of LMW-PTP activity. Full-length LMW-PTP shares 86% aa sequence identity with mouse LMW-PTP.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.