

DESCRIPTION

Species Reactivity	Human
Specificity	Detects CCL23/Ckβ8-1 in direct ELISAs and Western blots.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human CCL23/Ckβ8-1 Met46-Asn137 Accession # AAD10846
Conjugate	Alexa Fluor 350 Excitation Wavelength: 346 nm Emission Wavelength: 442 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Neutralization	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

CCL23 (SCY A23), a splice variant of CKβ8 (also known as MPIF-1), is a β chemokine isolated from the THP-1 cDNA library. CCL23 cDNA encodes a 137 amino acid (aa) residue precursor protein with a putative 21 aa residue signal peptide. Compared to CKβ8-1, CKβ8/MPIF-1 lacks a 17 aa residue stretch (Leu 47-Gly 63) present in CCL23 and has a unique arginine at residue 46. CKβ8-1/CKβ8 (MPIF-1) and leukotactin/MIP-1δ form a subgroup of β chemokines that have 6 conserved cysteine residues and an extended amino-terminus preceding the conserved cysteine pair. CCL23 shares approximately 73% nucleotide sequence identity with leukotactin (MIP-1δ), and the two genes have been localized to human chromosome 17 within 200 kb of each other. CCL23 mRNA expression has been detected in the pancreas, heart and skeletal muscle. Both CKβ8-1 and CKβ8 (MPIF-1) are potent agonists of CCR1 and have been shown to chemoattract peripheral blood lymphocytes and monocytes. The two chemokines have also been reported to chemoattract neutrophils and to inhibit colony formation by human hematopoietic progenitor cells. Similar to CKβ8 (MPIF-1), the truncated form of CKβ8-1 (aa 46-132) produced at R&D Systems has been shown to have greatly enhanced biological activity compared to the full length mature CKβ8-1 (aa 22-132).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.