

DESCRIPTION

Species Reactivity	Human/Mouse/Rat
Specificity	Detects endogenous human, mouse, and, rat PLC-β4 in Western blots.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human PLC-β4 Lys75-Gln250 Accession # Q15147
Conjugate	Alexa Fluor 594 Excitation Wavelength: 590 nm Emission Wavelength: 617 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

PLC-β4 (phospholipase C-beta 4; also PLCretA) is a 130 kDa member of the β-type PLC isozyme family. It is notably expressed in cerebellum and retina. PLC-β4 is associated with rhodopsin family GPCRs, where activation of Gq subunits initiates PLC-β4 hydrolysis of PIP2. Human PLC-β4 is 1175 amino acids (aa) in length. It contains a catalytic X-box (aa 313-463) and Y-box (aa 565-681), followed by a calcium-binding C2 domain (aa 688-786). There are multiple isoforms. One shows an alternate start site at Met114, a second contains a 40 aa substitution for the C-terminal 22 aa, a third shows a 14 aa substitution for the N-terminal 167 aa, and other forms demonstrate combinations of these alternative splice forms. Over aa 75-250, human PLC-β4 is 97% aa identical to mouse PLC-β4.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.