

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human CD99-L2 in direct ELISAs and Western blots.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant human CD99-L2 Val20-Ala188 Accession # Q8TCZ2
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

CD99 antigen-like 2 (CD99-L2) is a 45 kDa type I transmembrane glycoprotein in the CD99 family of molecules (1-3). The major form of human CD99-L2 cDNA encodes a 262 amino acid (aa) precursor with a 25 aa predicted signal sequence, a 160 aa extracellular domain (ECD), a 21 aa transmembrane (TM) segment, and a 56 aa cytoplasmic region (4). This form is called the long form, or isoform E3'-E4'-E3-E4. Other isoforms include the muscle E3-E4, missing aa 45-93, and E4, a short form missing aa 45-116 (1, 4). E3-E4 and E3'-E3-E4 forms are the major isoforms in mouse and rat, respectively (1). Human forms that diverge at Pro 45 (145 aa precursor) or Met 180 (173 aa precursor) have been sequenced, and would be predicted to lack a TM segment (4, 5). None of the forms contain predicted N-linked glycosylation sites within the ECD, but O-linked glycosylation is likely (1, 2). The ECD of the human CD99-L2 isoform E3-E4 shares 85%, 75% and 70% aa identity with the corresponding forms of mouse, rat, and bovine CD99-L2, respectively. The human CD99 and CD99-L2 ECDs share only about 35% aa identity, but both contain three conserved acidic motifs and are thought to originate from the same ancestral gene (1, 2). The nearly ubiquitous expression of CD99-L2 is similar to that of CD99. Human CD99-L2 cDNA is detected in most organs, but not in thymus (1). In the mouse, protein is detectable in lung, thymocytes, mouse leukocytes and vascular endothelial cells (1, 3, 6). The endothelial cell CD99-L2 is reported to mediate cell aggregation and neutrophil or monocyte, but not lymphocyte, extravasation to inflamed tissue *in vivo* (3, 6).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.