

DESCRIPTION

Species Reactivity	Rat
Specificity	Detects rat CXCL2/GROβ/MIP-2/CINC-3 in direct ELISAs and Western blots. In direct ELISAs and Western blots, approximately 20% cross-reactivity with recombinant rat (rr) CINC-2α is observed, 15% cross-reactivity with rrCINC-2β, recombinant h
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant rat CXCL2/GROβ/MIP-2/CINC-3 Ser32-Asn100 Accession # P30348
Conjugate	Alexa Fluor 350 Excitation Wavelength: 346 nm Emission Wavelength: 442 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Neutralization	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Cytokine-induced neutrophil chemoattractant 3 (CINC-3) has been found to be expressed by cytokine-stimulated rat alveolar macrophages and fibroblasts. Based on its protein and DNA sequences, CINC-3 (also known as rat MIP-2, GROβ and CXCL2) is a member of the alpha (CXC) subfamily of chemokines.

CINC-3 cDNA encodes a 100 amino acid (aa) residue precursor protein with a 31 aa signal peptide that is cleaved to yield a 69 aa mature secreted protein. The protein sequence of rat CINC-3 shares approximately 88% identity with murine MIP-2. Characteristic of ELR containing CXC chemokines, CINC-3 is known to be a potent chemotactic factor for rat neutrophils *in vitro* and *in vivo*. On the basis of cross-desensitization results of the various CINC proteins, it has been postulated that rat neutrophils have at least two classes of CINC receptors: a class of CINC-3 specific receptors as well as a second common receptor shared by all CINC3s.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.