

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human IL-34 in direct ELISAs and Western blots. In direct ELISAs, approximately 30% cross-reactivity with recombinant mouse IL-34 is observed.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant human IL-34 Asn21-Pro242 Accession # NP_689669
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Neutralization	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Interleukin 34 (IL-34; also known as uncharacterized protein C16orf77) is secreted as a homodimer consisting of 39 kDa monomers (1). It belongs to no known cytokine family. Human IL-34 is synthesized as a 242 amino acid (aa) precursor that contains a 20 aa signal sequence and a 222 aa mature chain. The mature chain contains one potential site of N-linked glycosylation. Human IL-34 is 71% identical to mouse IL-34 on the amino acid level (1). IL-34 is expressed in various tissues, including the heart, brain, liver, kidney, spleen, thymus, testes, ovary, small intestine, prostate, and colon, and is most abundant in the spleen (1). The receptor for IL-34 is colony-stimulating factor 1 receptor (CSF-1R) (1). IL-34 stimulates monocyte proliferation (1). In functional studies, IL-34, like CSF-1, the other ligand for CSF-1R, stimulated phosphorylation of extracellular signal-regulated kinase-1 and -2 (ERK1/2) in human monocytes (1). In addition, IL-34 promoted the formation of the colony-forming unit-macrophage (CFU-M), a macrophage progenitor, in human bone marrow cultures (1).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.