

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human GPR115 in direct ELISAs and Western blots. In direct ELISAs, less than 5% cross-reactivity with recombinant human (rh) GPR111, rhGPR114, and rhGPR125 is observed.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant human GPR115 Ser22-Ala347 Accession # Q8IZF3
Conjugate	Alexa Fluor 594 Excitation Wavelength: 590 nm Emission Wavelength: 617 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

GPR115 is a member of the LN-7TM family of adhesion-type 7-transmembrane (TM) G-protein coupled receptors (GPCR) that show a long extracellular N-terminus (1, 2). The 695 amino acid (aa) human GPR115 sequence predicts a 21 aa signal sequence, a 385 aa N-terminal extracellular domain (ECD), seven TM regions separated by 6-24 aa intracellular and extracellular regions, and a 40 aa cytoplasmic tail. Like other LN-7TM members, the ECD contains a highly glycosylated mucin-like stalk that is predicted to function in adhesion. This is followed by a cysteine-rich GPCR proteolytic cleavage site (GPS) (1). GPS domains, which have been described in other 7TM proteins including ETL, GPR126, HE6, and Latrophilin-1, are cleavage sites for processing proteins into two subunits (3-7). Within the N-terminal region that ends with the predicted cleavage site (aa 22-347), human GPR115 shares 58% aa sequence identity with the corresponding region of mouse and rat GPR115. GPR115 was identified from expressed sequence tags (ESTs) found in pregnant uterus, breast, and the genitourinary tract (1).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.