

DESCRIPTION


Species Reactivity	Human/Mouse
Specificity	Detects endogenous human/mouse WASF3/WAVE3 in Western blots.
Source	Polyclonal Goat IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli</i> -derived recombinant human WASF3/WAVE3 Met1-Lys180 Accession # Q9UPY6
Formulation	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose. See Certificate of Analysis for details. *Small pack size (-SP) is supplied either lyophilized or as a 0.2 µm filtered solution in PBS.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

	Recommended Concentration	Sample
Western Blot	1 µg/mL	See Below

DATA



PREPARATION AND STORAGE

Reconstitution	Reconstitute at 0.2 mg/mL in sterile PBS.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature recommended below. *Small pack size (-SP) is shipped with polar packs. Upon receipt, store it immediately at -20 to -70 °C
Stability & Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. <ul style="list-style-type: none"> 12 months from date of receipt, -20 to -70 °C as supplied. 1 month, 2 to 8 °C under sterile conditions after reconstitution. 6 months, -20 to -70 °C under sterile conditions after reconstitution.

BACKGROUND

WASF3/WAVE3 (Wiscott-Aldrich syndrome protein family Verprolin-homologous protein 3) is a 70 kDa member of the SCAR/WAVE family of proteins. It is expressed in select diverse cell types such as neurons, platelets, and breast epithelium. Upon activation, WASF3/WAVE3 is phosphorylated on Tyr151/248/337/486 and, in complex with HSPC300, initiates actin polymerization. WASF3/WAVE3 also regulates MMP expression. Human WASF3/WAVE3 is 502 amino acids (aa) in length. It contains one coiled-coil region (aa 57-93), three sequential poly-Pro regions (aa 304-410) and one VPH domain (aa 440-457) that binds to actin and Arp2/3. There is one isoform that shows a 36 aa substitution for aa 181-239. Over aa 1-180, human WASF3 shares 98% aa identity with mouse WASF3.