

Human α-Taxilin Alexa Fluor® 594-conjugated Antibody

Antigen Affinity-purified Polyclonal Sheep IgG Catalog Number: AF5575T

100 µg

DESCRIPTION	
Species Reactivity	Human
Specificity	Detects human α-Taxilin in direct ELISAs and Western blots.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	<i>E. coli-</i> derived recombinant human α-Taxilin Met1-Lys162 Accession # P40222
Conjugate	Alexa Fluor 594 Excitation Wavelength: 590 nm Emission Wavelength: 617 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet
	(SDS) for additional information and handling instructions.

APPLICATION

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

Immunocytochemistry Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE	
Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

 α -Taxilin (TXLN- α ; also p75 [in rat] and IL-14 α) is the founding member of the taxilin family of proteins. Although its predicted MW is 62 kDa, it runs anomalously at 74-76 kDa in SDS-PAGE. α -Taxilin is ubiquitously expressed and participates in intracellular vesicle trafficking. In particular, α -Taxilin interacts with syntaxin family members that localize to the plasma membrane, providing for SNARE complex formation. α -Taxilin also regulates the placement of growing protein chains. Human α -Taxilin is 546 amino acids (aa) in length. Its defining characteristic is a large coiled-coil region between aa 186-491. When an alternate start site at Met176 is used, a secreted cytokine is generated termed IL-14 α . Over aa 1-162, human α -Taxilin shares 81% aa identity with mouse α -Taxilin.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.

Rev. 9/16/2025 Page 1 of 1

Global | bio-techne.com info@bio-techne.com techsupport@bio-techne.com TEL: 1.612.379.2956

China | info.cn@bio-techne.com TEL: 400.821.3475