

DESCRIPTION

Species Reactivity	Rat
Specificity	Detects rat IL-7 Rα/CD127 in direct ELISAs and Western blots. In direct ELISAs, approximately 60% cross-reactivity with recombinant mouse IL-7 Rα and less than 10% cross-reactivity with recombinant human IL-7 Rα is observed.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant rat IL-7 Rα/CD127 Glu21-Asp239 Accession # NP_001099888
Conjugate	Alexa Fluor 488 Excitation Wavelength: 488 nm Emission Wavelength: 515-545 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

CyTOF-ready	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.
Flow Cytometry	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

IL-7 Rα (IL-7 receptor alpha; also CD127) is a 60-70 kDa member of the type I cytokine receptor family of molecules. It is expressed on resting naïve and memory CD8+ T cells, DN thymocytes, pre-B cells, Th1 CD4+ T cells, dendritic cells and monocytes. IL-7 Rα heterodimerizes with both the γc chain to create the IL-7 receptor, and the TSLPR subunit to generate the TSLP receptor. IL-7 Rα participates in T cell differentiation, naïve T cell survival, and CD4+ thymocyte proliferation. Mature rat IL-7 Rα is a 437 aa type I transmembrane glycoprotein. It contains a 219 aa extracellular region (aa 21-239) that possesses one type III fibronectin domain (aa 129-222), and a 193 aa cytoplasmic tail that shows a key phosphorylation site at Tyr447. Over aa 21-239, rat IL-7 Rα shares 79% and 67% aa identity with mouse and human IL-7 Rα, respectively.

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc., and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.