

DESCRIPTION

Species Reactivity	Mouse
Specificity	Detects mouse Siglec-1/CD169 in direct ELISAs and Western blots. In direct ELISAs, approximately 30% cross-reactivity with recombinant human (rh) Siglec-1 is observed and less than 1% cross-reactivity with rhSiglec-2 and recombinant mouse Siglec-1.
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	Mouse myeloma cell line NS0-derived recombinant mouse Siglec-1/CD169 Thr20-Leu1639 Accession # Q62230
Conjugate	Alexa Fluor 700 Excitation Wavelength: 675-700 nm Emission Wavelength: 723 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide
*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.	

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Neutralization	Optimal dilution of this antibody should be experimentally determined.
Western Blot	Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Siglecs are sialic acid specific I-type lectins that belong to the immunoglobulin superfamily. Structurally, they are transmembrane proteins with an N-terminal Ig-like V-set domain followed by varying numbers of Ig-like C2-set domains (1, 2). Mouse Siglec-1, also known as sialoadhesin and CD169, is a 175-185 kDa glycoprotein that consists of a 1619 amino acid (aa) extracellular domain (ECD) with one Ig-like V-set domain and 16 Ig-like C2-set domains, a 21 aa transmembrane segment, and a 35 aa cytoplasmic domain (3, 4). Within the ECD, mouse Siglec-1 shares 73% and 83% aa sequence identity with human and rat Siglec-1, respectively. Alternate splicing generates a soluble form of the ECD and a soluble isoform that is truncated following the first three Ig-like domains (3). Siglec-1 expression is restricted to lymph node and spleen macrophages and some tissue macrophages (4). The adhesive function of Siglec-1 is supported by the N-terminal Ig-like domain which shows a selectivity for α-2,3-linked sialic acid residues (4-6). Siglec-1 binds a number of sialylated molecules including the mannose receptor, MGL1, MUC1, PSGL-1, and different glycoforms of CD43 (7-10). Its binding capacity can be masked by endogenous sialylated molecules (11, 12). The sialylated and sulfated N-linked carbohydrates that modify Siglec-1 itself are required for ligand binding (7, 8). Siglec-1 is expressed on dendritic cells following rhinovirus exposure, and these DC promote T cell anergy (13). It is also induced on circulating monocytes during systemic sclerosis and HIV-1 infection (14-16). Siglec-1 can trap HIV-1 particles for trans-infection of permissive cells (15).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.