

DESCRIPTION

Species Reactivity	Human
Specificity	Detects human Olfactomedin-2/Noelin-2 in direct ELISAs and Western blots. In direct ELISAs, approximately 10% cross-reactivity with recombinant mouse (rm) Olfactomedin-1 is observed and less than 5% cross-reactivity with recombinant hu
Source	Polyclonal Sheep IgG
Purification	Antigen Affinity-purified
Immunogen	Chinese hamster ovary cell line CHO-derived recombinant human Olfactomedin-2/Noelin-2 Gln21-Pro454 Accession # O95897
Conjugate	Alexa Fluor 532 Excitation Wavelength: 534 nm Emission Wavelength: 553 nm
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide *Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.

APPLICATIONS

Please Note: Optimal dilutions should be determined by each laboratory for each application. [General Protocols](#) are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

PREPARATION AND STORAGE

Shipping The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.

Stability & Storage Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied

BACKGROUND

Olfactomedin-2 (OLFM2, OM2), also called OlfC or neuronal olfactomedin-related endoplasmic reticulum-localized-2 (Noelin-2) is a 50 kDa (predicted), secreted member of the olfactomedin/noelin family of proteins (1-3). Since Noelin-2 has also been used to designate an alternately spliced form of Noelin-1, Olfactomedin-2 is the less ambiguous name (4). Olfactomedin domain-containing proteins are often found in neural tissues (3). Human Olfactomedin-2 expressed sequence tags have mainly been found in the brain and eye, and RNA in the cerebellum (1, 2). In zebrafish, Olfactomedin-2 expression is developmentally regulated and, like Noelin-1, may play a role in neural crest cell differentiation (5). The human Olfactomedin-2 cDNA encodes a 20 amino acid (aa) signal sequence and a 454 mature protein that contains two coiled-coil regions (aa 58-85 and 136-193) and an Olfactomedin domain (aa 194-446). Mature human Olfactomedin-2 shares 98% aa identity with rat, bovine and equine, 97% with mouse and canine, 81% with *Xenopus*, and 77% with zebrafish Olfactomedin-2. No alternate splice variants have been reported for Olfactomedin-2 (1). Polymorphisms of human Olfactomedin-2, with or without additional polymorphisms in optineurin (OPTN), have been associated with a small fraction of high intraocular pressure or open-angle glaucoma cases in Japanese patients (6). The family member myocilin shows stronger association with these ocular disorders, and either myocilin or Noelin-1 might heterodimerize with Olfactomedin-2 (1). Evidence suggests that Olfactomedin-2 has evolved from Noelin-1 gene duplication, and myocilin from Olfactomedin-2 gene duplication (1).

PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.