

## Human PLD1 Alexa Fluor® 750-conjugated Antibody

Antigen Affinity-purified Polyclonal Goat IgG Catalog Number: AF5615S

100 µg

DESCRIPTION				
Species Reactivity Human				
Specificity	Detects endogenous human PLD1 in Western blots.			
Source	Polyclonal Goat IgG			
Purification	Antigen Affinity-purified			
Immunogen	E. coli-derived recombinant human PLD1 Met1-Pro140 Accession # Q13393			
Conjugate	Alexa Fluor 750 Excitation Wavelength: 749 nm Emission Wavelength: 775 nm			
Formulation	Supplied 0.2mg/ml in 1X PBS with RDF1 and 0.09% Sodium Azide			
	*Contains <0.1% Sodium Azide, which is not hazardous at this concentration according to GHS classifications. Refer to the Safety Data Sheet (SDS) for additional information and handling instructions.			

## **APPLICATIONS**

Please Note: Optimal dilutions should be determined by each laboratory for each application. General Protocols are available in the Technical Information section on our website.

Western Blot Optimal dilution of this antibody should be experimentally determined.

China | info.cn@bio-techne.com TEL: 400.821.3475

PI	REP	'ARA	ATIO	N AI	ND.	STO	RA	GΕ

Shipping	The product is shipped with polar packs. Upon receipt, store it immediately at the temperature recommended below.		
Stability & Storage	Protect from light. Do not freeze. 12 months from date of receipt, 2 to 8 °C as supplied		

## **BACKGROUND**

PLD1 (Phospholipase D1a) is a 110-120 kDa member of the phospholipase D family of enzymes. It is expressed in endothelial cells and select tissues. Following activation and association with Type Iα PIPkinase, PLD1 hydrolyzes the phosphodiester bond of membrane phosphatidylcholine, generating phosphatidic acid. Human PLD1 is 1074 amino acids in length). It contains one PX/phox homology domain (aa 84-206), a pleckstrin homology domain (aa 219-328), and two phosphodiesterase enzyme regions (aa 459-486 and 891-918). There are three splice variants. One shows a 10 aa substitution for aa 962-1074 (PLD1d), a second shows an 84 aa substitution for aa 514-1074 (PLD1c), and a third shows an Asn substitution for aa 585-623 (PLD1b). Caspase cleavage of PLD1a at Gln545-Ser546 generates a 62 kDa N-terminal, and a 61 kDa C-terminal fragment. Over aa 1-140, human PLD1a shares 91% aa identity with mouse PLD1a.

## PRODUCT SPECIFIC NOTICES

This product is provided under an agreement between Life Technologies Corporation and R&D Systems, Inc, and the manufacture, use, sale or import of this product is subject to one or more US patents and corresponding non-US equivalents, owned by Life Technologies Corporation and its affiliates. The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components (1) in manufacturing; (2) to provide a service, information, or data to an unaffiliated third party for payment; (3) for therapeutic, diagnostic or prophylactic purposes; (4) to resell, sell, or otherwise transfer this product or its components to any third party, or for any other commercial purpose. Life Technologies Corporation will not assert a claim against the buyer of the infringement of the above patents based on the manufacture, use or sale of a commercial product developed in research by the buyer in which this product or its components was employed, provided that neither this product nor any of its components was used in the manufacture of such product. For information on purchasing a license to this product for purposes other than research, contact Life Technologies Corporation, Cell Analysis Business Unit, Business Development, 29851 Willow Creek Road, Eugene, OR 97402, Tel: (541) 465-8300. Fax: (541) 335-0354.

Rev. 9/16/2025 Page 1 of 1

Global | bio-techne.com info@bio-techne.com techsupport@bio-techne.com TEL: 1.612.379.2956